

Torsion primes in loop space homology

Y. Felix, S. Halperin and J.-C. Thomas

February 1, 2008

Abstract

We construct a finite 1-connected CW complex X such that $H_*(\Omega X; \mathbb{Z})$ has p -torsion for the infinitely many primes satisfying $p \equiv 5, 7, 17, 19 \pmod{24}$, but no p -torsion for the infinitely many primes satisfying $p \equiv 13$ or $23 \pmod{24}$.

AMS Classification : 55P35

Key words : loop space homology

A *homology torsion prime* for a topological space Y is a prime p such that $H_*(Y; \mathbb{Z})$ has p torsion. In [3] and [4] respectively, D. Anick and L. Avramov constructed simply connected finite CW complexes X of dimension 4 such that $H_*(\Omega X, \mathbb{Z})$ has p -torsion for all primes p . Since there are only countably many homotopy types of simply connected finite CW complexes, this immediately raised the

Question. If $\mathcal{P}$ denotes the set of all primes, what are the conditions for a subset $\mathcal{D} \subset \mathcal{P}$, to be the set of homology torsion primes of the loop space of a finite simply connected CW complex ?

Of course, any finite set can be realized. It suffices to take a product of Moore spaces. On the other hand by a result of McGibbon and Wilkerson [7], if X is rationally elliptic (i.e., $\dim \pi_*(X) \otimes \mathbb{Q} < \infty$), then the set of homology torsion primes for ΩX is finite.

In this note we adapt the construction of Anick to prove

Theorem 1. *There exists a 4-dimensional finite simply connected CW complex X such that the homology torsion primes for ΩX are an infinite set in $\mathcal{P}$, with an infinite complement.*

Theorem 2. *For each finite subset E of $\mathcal{P}$ there exists a 4-dimensional finite simply connected CW complex X_E which has $\mathcal{P} \setminus E$ as the set of homology torsion primes for ΩX_E .*

Our examples are CW complexes X obtained by attaching thirteen 4-cells to a wedge of eight 2-spheres via a map $f : Y = \bigvee_{i=1}^{13} S_i^3 \rightarrow Z = \bigvee_{j=1}^8 S_j^2$. We label the 2-spheres $u_1, u_2, u_3, u_4, v, w, x_1, x_2$ and, fixing six integers a, b, c, d, a_2, b_2 , attach the 13 cells in degree 4 along the elements

$$\begin{aligned} &[x_1, u_1] - [u_1, v] - a[u_2, v], \quad [x_1, u_2] - b[u_2, v] - d[u_3, v], \quad [x_1, u_3] - c[u_2, v] \\ &[x_1, u_4], \quad [x_1, v], \quad [x_1, w], \quad [x_2, u_1], \quad [x_2, u_3], \quad [x_2, u_4] \\ &[x_2, v], \quad [x_2, w], \quad [x_2, u_2] - [u_4, v], \quad [u_1, w] + a_2[u_2, w] + b_2[u_3, w]. \end{aligned}$$

Denote by $R\langle x_1, \dots, x_n \rangle$ the tensor algebra over a commutative ring R with generators $x_1, \dots, x_n$. The image A_X of $H_*(\Omega Z; \mathbb{Z})$ in $H_*(\Omega X; \mathbb{Z})$ is the quotient $H_*(\Omega Z; \mathbb{Z})/I$,

where I denotes the two-sided ideal generated by the image of $H_2(\Omega f)$ ([6]). Thus $A_X = \mathbb{Z}\langle u_1, u_2, u_3, u_4, v, w, x_1, x_2 \rangle / I$, where all variables have degree one and I is the ideal generated by the relations above. For any field k , the Hilbert series of $A_X \otimes k$ and of $H_*(\Omega X; k)$ are related by a formula of Roos ([8]; Roos asserts this explicitly only for $k = \mathbb{Q}$, but as Avramov observed in [4] it holds for any coefficient field)

$$P_{\Omega X}(t)^{-1} = (1+t)A(t)^{-1} - t + 8t^2 - 13t^3.$$

In particular $H_*(\Omega X, \mathbb{Z})$ and A_X have the same torsion primes.

Our next step is to construct a graded algebra E and then to identify the algebra A_X as the semi-tensor product ([9]) of the tensor algebra $\mathbb{Z}\langle x_1, x_2 \rangle$ with E . Put $F = \mathbb{Z}\langle u_1, u_2, u_3, u_4, v, w \rangle$. For $i = 1, \dots, 4$, define elements $\sigma_{i,m}, \rho_{i,m} \in F$ by $\sigma_{i,1} = u_i$, $\sigma_{i,m+1} = [\sigma_{i,m}, v]$ and $\rho_{i,m} = [\sigma_{i,m-1}, w]$. Then define integers a_m and b_m , $m \geq 3$, by

$$\begin{pmatrix} a_m \\ b_m \end{pmatrix} = \begin{pmatrix} a \\ 0 \end{pmatrix} + \begin{pmatrix} b & c \\ d & 0 \end{pmatrix} \begin{pmatrix} a_{m-1} \\ b_{m-1} \end{pmatrix}.$$

Finally, set

$$\tau_m = \rho_{1,m} + a_m \rho_{2,m} + b_m \rho_{3,m}, \quad m \geq 2$$

and set

$$S = \{\tau_m, m \geq 2\} \cup \{a_m \rho_{4,m+1}, m \geq 2\}.$$

Define E to be the quotient of F by the two-sided ideal J generated by S .

A unique action of x_1 and x_2 by derivations in F is given by

$$\begin{cases} x_1 * u_1 = [u_1, v] + a[u_2, v] \\ x_1 * u_2 = b[u_2, v] + d[u_3, v] \\ x_1 * u_3 = c[u_2, v] \\ x_1 * u_4 = x_1 * v = x_1 * w = 0 \\ x_2 * u_1 = x_2 * u_3 = x_2 * u_4 = x_2 * v = x_2 * w = 0 \\ x_2 * u_2 = [u_4, v] \end{cases}$$

This extends uniquely to an action of $\mathbb{Z}\langle x_1, x_2 \rangle$ in F . The action of $\mathbb{Z}\langle x_1, x_2 \rangle$ preserves J , and so an action is induced in $F/J = E$. Moreover, a presentation of the semi-tensor product of $\mathbb{Z}\langle x_1, x_2 \rangle$ with E is given by

$$\mathbb{Z}\langle x_1, x_2 \rangle \otimes E = \mathbb{Z}\langle u_1, u_2, u_3, u_4, x_1, x_2, v, w \rangle / I,$$

where I is the two-sided ideal generated by the relations $[x_i, u_j] = x_i * u_j$, $[x_i, v] = x_i * v$, $[x_i, w] = x_i * w$ and $[u_1, w] + a_2[u_2, w] + b_2[u_3, w] = 0$. This identifies the semi-tensor product with A_X and shows that the torsion primes for A_X are exactly those of E .

By construction, for $m \geq 3$, the elements $\rho_{4,m}$ are torsion elements in A_X . In the same way as in [3], we have

Lemma 1. *If b is an integer, then $b \cdot \rho_{4,m} \neq 0$ in A_X unless $b \equiv 0 \pmod{a_{m-1}}$.*

Therefore all the prime numbers that appear as divisor of one of the a_m are torsion primes for A_X . The next lemma shows that all the torsion primes are of that form.

Lemma 2. *If p does not divide one of the a_m , $m \geq 2$, then A_X has no element of order p .*

proof: Invert the non-zero integers a_m , $m \geq 2$ and set $R = \mathbb{Z}(\frac{1}{a_m})_{m \geq 2}$. It is immediate from the criteria in [2] that the elements $\rho_{4,m+1}$, τ_m , $m \geq 2$, form a strongly free (or inert) sequence in $F \otimes R$, and it follows that $E \otimes R = (F \otimes R)/J$ splits back as a sub R -module of $F \otimes R$. In particular, it is torsion free. Hence so is $A_X \otimes R$. $\square$

Proof of Theorem 1. Consider the particular case where $a = 0, b = 4, c = -3, d = 1, a_2 = 11, b_2 = 5$. In that case

$$\tau_m = \rho_{1,m} + (2 + 3^m)\rho_{2,m} + (2 + 3^{m-1})\rho_{3,m}$$

The integer p is a homology torsion prime for ΩX if and only if for some $m \geq 2$, $2 + 3^m \equiv 0 \pmod p$.

We show now that the equation $2 + 3^m \equiv 0 \pmod p$ has a solution when $p \equiv 5, 7, 17$ or $19 \pmod{24}$, and does not have a solution when $p \equiv 13$ or $23 \pmod{24}$.

In fact, recall that a number a is a quadratic residue mod p if and only if the Legendre symbol $(a/p) = 1$. Recall ([5], p. 209, and Theorem 9-10), that

$$(3/p) = \begin{cases} 1 & \text{if } p \equiv 1 \text{ or } 11 \pmod{12} \\ -1 & \text{if } p \equiv 5 \text{ or } 7 \pmod{12} \end{cases}$$

$$(-2/p) = \begin{cases} 1 & \text{if } p \equiv 1 \text{ or } 3 \pmod{8} \\ -1 & \text{if } p \equiv 5 \text{ or } 7 \pmod{8} \end{cases}$$

Therefore if $p \equiv 5, 7, 17$ or $19 \pmod{24}$, the integer 3 is not a quadratic residue and is therefore a primitive root mod p . Therefore the equation $2 + 3^m \equiv 0 \pmod p$ has always a solution. On the other hand when $p \equiv 13$ or $23 \pmod{24}$, then 3 is a quadratic residue, but -2 is not, and the equation $-2 \equiv 3^n \pmod p$ has therefore no solution.

Recall now Dirichlet's Theorem ([5], Theorem 3-7): When r and s are relatively prime integers there are infinitely many primes q such that $q \equiv r \pmod s$. In particular there are infinitely many prime numbers p such that $p \equiv 5 \pmod{24}$ and infinitely many prime numbers such that $p \equiv 13 \pmod{24}$. This shows that the set of torsion primes for A_X is an infinite subset of $\mathcal{P}$ with an infinite complement. $\square$

Proof of Theorem 2. Consider the particular case where $b = 1, c = 0, d = 0, a_2 = 1, b_2 = 0$. In that case $a_m = 1 + a(m - 2)$. Thus $a_m \equiv 0 \pmod p$ for some m if and only if $a \not\equiv 0 \pmod p$. Let $\mathcal{S} = \{p_1, \dots, p_n\}$ be any finite subset of $\mathcal{P}$ and put $a = p_1 p_2 \cdots p_n$. Then p is a prime divisor for A_X if and only if $p \notin \mathcal{S}$. $\square$

References

- [1] J.F. Adams and P.J. Hilton, On the chain algebra of a loop space, *Comment. Math. Helv.* **30** (1956), 305-330.
- [2] D. Anick, Non-commutative graded algebras and their Hilbert series, *J. of Algebra*, **78** (1982), 120-140.
- [3] D. Anick, A loop space whose homology has torsion of all orders, *Pacific Journal of Mathematics*, **123** (1986), 257-262.
- [4] L. Avramov, Torsion in loop space homology, *Topology* **25** (1986), 155-157.
- [5] D. Burton, *Elementary number theory*, Allyn and Bacon, 1980.
- [6] J.-M. Lemaire, *Algèbres connexes et homologie des espaces de lacets*, Lecture Notes in Math **422**, Springer-Verlag, 1972.
- [7] C.A. McGibbon and C.W. Wilkerson, Loop spaces of finite complexes at large primes, *Proc. Amer. Math. Soc.* **96** (1986), 698-702.
- [8] J.E. Roos, Relations between the Poincaré-Betti series of loop spaces and of local rings, *Lecture Notes in Math.* **740**, Springer-Verlag, Berlin 1979, pp. 285-322.

- [9] L. Smith, Split extensions of Hopf algebras and semi-tensor products, *Math. Scand.* **26** (1970), 17-41.

Institut de Mathématiques,
Université Catholique de Louvain
1348 Louvain-La-Neuve, Belgium

College of Computer, Mathematical and Physical Sciences
University of Maryland
College Park, MD 20742-3281, USA

Faculté des Sciences
Université d'Angers
Bd. Lavoisier, 49045 Angers, France